


ATTACHMENT 1




If nervousness, dizziness or sleeplessness occur, discontinue use and consult a doctor. Do not take this product if you have heart disease, high blood pressure, thyroid disease, diabetes or difficulty in urination due to enlargement of the prostate gland unless directed by a doctor. Do not exceed recommended dosage. Keep this and all drugs out of the reach of children. In case of accidental overdose, contact a doctor or poison control center immediately. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. As with any drug, if you are pregnant or nursing a baby, seek the advice of a health professional before using this product. Do not use with other products containing acetaminophen. **Drug Interaction Precaution:** Do not use this product if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you are uncertain whether your prescription drug contains an MAOI, consult a health professional before taking this product. **Directions:** Adults and children 12 years of age and older: Take 2 caplets every 4-6 hours. Do not take more than 8 caplets in 24 hours, or as directed by a doctor. Children under 12 years: Do not use this adult product in children under 12 years of age. This will provide more than the recommended dose (overdose) and could cause serious health problems. **Inactive Ingredients:** Carnauba Wax, Cellulose, Corn Starch, D&C Yellow 10, FD&C Blue 1, FD&C Red 40, Hydroxypropyl Methylcellulose, Iron Oxide, Magnesium Stearate, Polyethylene Glycol, Polysorbate 80, Sodium Starch Glycolate, Titanium Dioxide. Do not use if pouch is opened. Store at room temperature.

MCNEIL-PPC, INC., FORT WASHINGTON, PA 19034 USA
© MCN-PPC, Inc. '01

5331160

150% OF SIZE



If nervousness, dizziness or sleeplessness occur, discontinue use and consult a doctor. Do not take this product if you have heart disease, high blood pressure, thyroid disease, diabetes or difficulty in urination due to enlargement of the prostate gland unless directed by a doctor. Do not exceed recommended dosage. Keep this and all drugs out of the reach of children. In case of accidental overdose, contact a doctor or poison control center immediately. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. As with any drug, if you are pregnant or nursing a baby, seek the advice of a health professional before using this product. Do not use with other products containing acetaminophen. **Drug Interaction Precaution:** Do not use this product if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you are uncertain whether your prescription drug contains an MAOI, consult a health professional before taking this product. **Directions:** Adults and children 12 years of age and older: Take 2 caplets every 4-6 hours. Do not take more than 8 caplets in 24 hours, or as directed by a doctor. Children under 12 years: Do not use this adult product in children under 12 years of age. This will provide more than the recommended dose (overdose) and could cause serious health problems. **Inactive Ingredients:** Carnauba Wax, Cellulose, Corn Starch, D&C Yellow 10, FD&C Blue 1, FD&C Red 40, Hydroxypropyl Methylcellulose, Iron Oxide, Magnesium Stearate, Polyethylene Glycol, Polysorbate 80, Sodium Starch Glycolate, Titanium Dioxide. Do not use if pouch is opened. Store at room temperature.

MCNEIL-PPC, INC., FORT WASHINGTON, PA 19034 USA

© MCN-PPC, Inc. '01

5331160